

Properties of superdeformed fission isomers in the cranked relativistic Hartree-Bogoliubov theory.

A. V. Afanasjev^{1,a} and P. Ring²

¹ Physik-Department der Technischen Universität München,
D-85747 Garching, Germany

Received 1 January 1996; revised version 1 January 1997

Abstract. The rotational and deformation properties of superdeformed fission isomers in the $A \sim 240$ mass region have been investigated within the framework of the cranked relativistic Hartree-Bogoliubov theory. The dependence of the results of the calculations on the parametrization of the RMF Lagrangian has been studied. The rotational properties are best described by the NL1 force.

The region of $A \sim 240$ is the first one where the superdeformed (SD) shapes have been discovered experimentally [1] in the fission isomers in 1962. According to the measurements of the moments of inertia and the quadrupole moments they are shape isomers with the deformation much larger than the usual ground state deformation. Despite the fact that during almost 40 years numerous experimental efforts have been made for the study of fission isomers, the experimental data on their properties in some respects is more limited than the one collected in the last 15 years in the regions of superdeformation at high spin such as $A \sim 60, 80, 130, 150$ and 190 [2]. Recent advances in the experimental techniques leading to the observation of hyperdeformed (third) minimum [3–5] and vibrational structures in the SD (second) minimum [6–8] revived the interest to this mass region of superdeformation.

A number of theoretical investigations has been performed in this mass region studying different properties of fission isomers within the frameworks of the phenomenological macroscopic + microscopic method and non-relativistic microscopic theories based on the zero-range Skyrme forces and finite range Gogny forces, see Refs. [9–13] and references therein. Recently, some properties of fission isomers in ^{226}Ra , ^{232}Th and ^{240}Pu have been studied in the relativistic mean field (RMF) theory by the Frankfurt group [14, 15]. These investigations, however, have been performed in the BCS approximation using a schematic treatment of pairing in the constant gap approximation with the pairing parameters adjusted to the experimental data in the first minimum.

The present investigation is aimed on a more systematic study of the properties

of SD fission isomers within the framework of RMF theory in the present state of the art. The results of this study will be presented in a forthcoming article [16]. In the present contribution, we will concentrate on the even-even nuclei in which rotational structures have been observed so far [17], namely on $^{236,238}\text{U}$ and ^{240}Pu nuclei.

Table 1. Kinematic moments of inertia ($J^{(1)}$), charge quadrupole (Q_0) and mass hexadecupole (Q_{40}) moments of SD band in ^{240}Pu calculated at rotational frequency $\Omega_x = 0.01$ MeV with different parametrizations of the RMF Lagrangian. In the case of the NL1 force, the results of the CRHB calculations with and without APNP(LN) [marked as 'LN' and 'Unpr'] and the results of the CRMF calculations with no pairing [marked as 'CRMF'] are given. For other forces, only the results of the CRHB calculations with APNP(LN) are shown.

Quantity	NLSH	NL3	NL1 (LN/Unpr/CRMF)	NL-Z	Exp.
$J^{(1)}$ [MeV $^{-1}$]	138.2	144.4	155.2 / 177.2 / 189.1	157.3	149.3
Q_0 [eb]	33.0	35.5	38.2 / 39.6 / 39.5	39.7	
Q_{40} [fm 4]	36121	41901	49425 / 56329 / 55533	53328	

As a theoretical tool we are using the recently developed Cranked Relativistic Hartree-Bogoliubov (CRHB) theory [18] which has been very successful in the description of the properties of SD bands in the $A \sim 190$ mass region [19, 18] and rare-earth nuclei [20]. Compared with previous relativistic studies in the $A \sim 240$ mass region, it has the following advantages: **(i)** the cranked RMF equations are solved on the Hartree-Bogoliubov level, **(ii)** the finite range D1S Gogny force is used in the particle-particle channel, **(iii)** approximate particle number projection is performed by means of the Lipkin-Nogami method (further APNP(LN)), **(iv)** the cranking model approximation is employed which allows to study the rotational properties of fission isomers and **(v)** this theory is formulated in three-dimensional Cartesian coordinates which allows to study the possible appearance of triaxial deformation. Since this study is concerned with the properties of the states in the second minimum, only reflection symmetric shapes are considered. This approximation is justified since octupole deformation becomes important only after the second well (see for example Refs. [11, 14]).

The calculated moments of inertia of the SD isomers in ^{236}U and ^{240}Pu are shown in Table 1 and Fig. 1. The change of the force from NLSH [21] via NL3 [22] and NL1 [23] to NL-Z [24] leads to the increase of the kinematic moments of inertia, charge quadrupole and mass hexadecupole moments (Table 1). The experimental moment of inertia of the SD band in ^{240}Pu [17] is located between the results of the calculations with NL3 and NL1 (see Table 1). The use of the NLSH and NL-Z forces for the RMF Lagrangian leads to larger deviations from experiment. The results of the calculations with NL3 underestimate the experimental moments of inertia by 3 – 5%. This is also the case for ^{238}U where experimental and calculated (with

NL3) moments of inertia are 153.1 and 142.8 MeV⁻¹. The difference between $J^{(1)}$ moments of inertia of ^{236,238}U nuclei is 3.8 MeV⁻¹ in experiment while it is only 0.6 MeV⁻¹ in the calculations with NL3. The NL1 force describes the absolute values of the moments of inertia and their difference in ^{236,238}U better than NL3. In these nuclei, the $J^{(1)}$ values calculated at $\Omega_x = 0.01$ MeV are 150.5 and 154.7 MeV⁻¹ which agree very well with the experimental values of 149.25 and 153.06 MeV⁻¹. On the other hand, the calculations with NL1 somewhat (by $\sim 4\%$) overestimate the experimental moments of inertia in ²⁴⁰Pu (see Fig. 1 and Table 1).




Fig. 1. Experimental and calculated kinematic ($J^{(1)}$) and dynamic ($J^{(2)}$) moments of inertia of SD fission isomers in ²³⁶U and ²⁴⁰Pu. The notation of the lines and symbols is given in the figure.

The CRHB calculations indicate that kinematic and dynamic moments of inertia increase with increasing rotational frequency Ω_x (Fig. 1). In addition, the difference between these moments grows with the increase of Ω_x . The experimental data in ²⁴⁰Pu shows such features, while they are not seen in ²³⁶U. These features are predominantly due to the gradual alignment of the $N = 8$ neutrons and $N = 7$ protons and a smooth decrease of pairing correlations with increasing Ω_x . They are similar to the ones observed in the $A \sim 190$ region of superdeformation, see Refs. [19, 18] and references quoted therein.

The importance of APNP(LN) for the description of the rotational and deformation properties of SD fission isomers is clearly seen on the example of the calculations performed with the NL1 force (Table 1). The CRMF calculations [25]

with no pairing give $J^{(1)} = 189.1 \text{ MeV}^{-1}$ which is below the rigid body moment of inertia ($J_{rig} = 211.7 \text{ MeV}^{-1}$) defined from the density distribution. The inclusion of pairing somewhat decreases the calculated moment of inertia which still considerably exceeds the experimental value. Only the CRHB results with APNP(LN) come close to experiment. One should also note that in the case of ^{240}Pu the inclusion of pairing without APNP(LN) has only a marginal effect on the charge quadrupole and mass hexadecupole moments of inertia (Table 1). On the contrary, APNP(LN) has a strong impact on these moments decreasing their values. In addition, APNP(LN) increases the strength of pairing correlations especially for the neutron subsystem. In the CRHB theory, the pairing energies are defined as $E_{pairing} = -\frac{1}{2}Tr(\Delta\kappa)$ [18]. The values of neutron and proton pairing energies obtained in the CRHB calculations without and with APNP(LN) are $E_{pairing}^\nu = -1.823 \text{ MeV}$, $E_{pairing}^\pi = -10.030 \text{ MeV}$ and $E_{pairing}^\nu = -13.129 \text{ MeV}$, $E_{pairing}^\pi = -13.603 \text{ MeV}$, respectively.

Table 2. Experimental and theoretical charge quadrupole moments of SD fission isomers. The results of the calculations with the NL1 and NL3 forces are given. Experimental data for U and Pu isotopes are taken from Ref. [17], while the one for ^{242}Am from Ref. [26].

	^{236}U	^{238}U	^{236}Pu	^{239}Pu	^{240}Pu	^{242}Am
$Q_0^{exp} \text{ (eb)}$	32 ± 5	29 ± 3	37 ± 10	36 ± 4		$35.5 \pm 1.0_{st} \pm 1.2_{mod}$
$Q_0^{NL1} \text{ (eb)}$	35.9	37.0	36.6		38.2	
$Q_0^{NL3} \text{ (eb)}$	33.8	34.0	35.0		35.5	

The charge quadrupole moments calculated with the forces NL1 and NL3 are compared with available experimental data in Table 2. One should note that the small error bars on the experimental values of Q_0 given for ^{238}U and ^{240}Am nuclei should be treated with caution since even modern experiments do not provide an accuracy of the absolute Q_0 values better than 15%, see discussion in Ref. [19]. In addition, when comparing the calculations with experiment one should take into account that (i) the Q_0^{exp} values have been obtained with different experimental techniques [17], (ii) it is reasonable to expect that an addition of one neutron to ^{239}Pu will not change considerably the Q_0 value and thus $Q_0^{exp}(^{239}\text{Pu})$ could be used for comparison with the calculated $Q_0(^{240}\text{Pu})$. With these considerations in mind, it is clear that the results of the calculations come reasonably close to experiment.

In conclusion, the cranked relativistic Hartree-Bogoliubov theory has been applied for the description of the rotational and deformation properties of superdeformed fission isomers in the $A \sim 240$ mass region. This theory does not employ any adjustable parameters and for the description of pairing correlations uses the well established D1S Gogny force. The study of the dependence of the results on the parametrization of the RMF Lagrangian reveals that rotational properties are best described by the NL1 force, while the uncertainties of the experimental data on charge quadrupole moments do not allow to use this experimental quantity for the selection of the best RMF force.

A.V.A. acknowledges support from the Alexander von Humboldt Foundation. This work is also supported in part by the Bundesministerium für Bildung und Forschung under the project 06 TM 979.

- a.* Alexander von Humboldt fellow, on leave of absence from the Laboratory of Radiation Physics, Institute of Solid State Physics, University of Latvia, LV 2169 Salaspils, Miera str. 31, Latvia

References

1. S. M. Polikanov *et al.*, *Sov. Phys. JETP* **15** (1962) 1016.
2. B. Singh, R. B. Firestone and S. Y. F. Chu, *Nuclear Data Sheets* **78** (1996) 1.
3. A. Krasznahorkay *et al.*, *Phys. Rev. Lett.* **80** (1998) 2073.
4. A. Krasznahorkay *et al.*, to be published
5. A. Krasznahorkay *et al.*, *Acta Phys. Pol.* **B30** (1999) 1467.
6. M. Hunyadi *et al.*, *Acta Phys. Pol.* **B30** (1999) 1263.
7. M. Hunyadi *et al.*, to be published
8. P. G. Thirolf, contribution to these proceedings
9. I. Hamamoto and W. Ogle, *Nucl. Phys.* **A240** (1975) 54.
10. J. Dudek, W. Nazarewicz and A. Faessler, *Nucl. Phys.* **A412** (1984) 61.
11. S. Ówiok, W. Nazarewicz, J. X. Saladin, W. Płóciennik and A. Johnson, *Phys. Lett.* **B322** (1994) 304.
12. J. F. Berger, M. Girod and D. Gogny, *Nucl. Phys.* **A 502** (1989) 85c.
13. S. J. Krieger, P. Bonche, H. Flocard, P. H. Heenen and M. S. Weiss, *Nucl. Phys.* **A 572** (1994) 384.
14. K. Rutz, J. A. Maruhn, P.-G. Reinhard and W. Greiner, *Nucl. Phys.* **A 590** (1995) 680.
15. M. Bender, K. Rutz, P.-G. Reinhard and J. A. Maruhn, *Eur. Phys. J. A*, in press (see also report nucl-th/9910025)
16. A. V. Afanasjev and P. Ring, in preparation.
17. D. Habs, *Nucl. Phys.* **A502** (1989) 105c.
18. A. V. Afanasjev, P. Ring and J. König, *Nucl. Phys. A*, in press (see also report nucl-th/0001054).
19. A. V. Afanasjev, J. König and P. Ring, *Phys. Rev.* **C60** (1999) 051303.
20. A. V. Afanasjev, J. König, P. Ring, L. M. Robledo and J. L. Egido, to be published.
21. M. M. Sharma, M. A. Nagarajan and P. Ring, *Phys. Lett.* **B312** (1993) 377.
22. G. A. Lalazissis, J. König and P. Ring, *Phys. Rev.* **C55** (1997) 540.
23. P.-G. Reinhard, M. Rufa, J. Maruhn, W. Greiner and J. Friedrich, *Z. Phys.* **A323** (1986) 13.
24. M. Rufa *et al.*, *Phys. Rev.* **C38** (1988) 390.
25. A. V. Afanasjev, J. König and P. Ring, *Nucl. Phys.* **A608** (1996) 107.
26. H. Backe *et al.*, *Phys. Rev. Lett.* **80** (1998) 920.